

LABORATORY TEST REPORT

Name	Md. ZAKIR HUSSAIN		
Sample ID	A1356837	Reg. No	0522411200018
Age/Gender	62 Years/Male	SPP Code	SPL-BH-144
Referred by	Dr. ALOK KUMAR MD.	Collected On	20-Nov-2024 08:22 AM
Referring Customer	Daffodils Laboratory	Received On	21-Nov-2024 09:41 AM
Primary Sample	Whole Blood	Reported On	23-Nov-2024 11:53 AM
Sample Tested In	Lithium Heparin	Report Status	Final Report
Client Address	New Millat Colony, MOB-8271440097		

IMMUNOLOGY & SEROLOGY

Test Name	Results	Units	Biological Reference Interval
QuantiFeron TB Gold (Interferon Gamma Release Assay)			
Gamma Interferon Nil Tube	0.26		
Gamma Interferon Antigen	1.99		
TB IGRA (IFN-γ) Levels <small>(Method: ELISA)</small>	1.73	IU/mL	Negative: < 0.35 Positive: >= 0.35
TB IGRA (IFN-γ) Results <small>(Method: ELISA)</small>	Positive		

Interpretation :

Nil (pg/mL)	TB Antigen minus Nil (pg/mL)	Results	Report/ Interpretation
<= 8.0	>= 0.35 and >=25 % N	Positive	M. tuberculosis infection likely
	< 0.35	Negative	M. tuberculosis infection not likely
	>= 0.35 and <=25 % N	Negative	M. tuberculosis infection not likely
> 8.0	Any Value	Indeterminate	Results are indeterminate for Antigen responsiveness

- A Negative QFT result does not preclude the possibility of M.tuberculosis infection or tuberculosis disease. False negative can be due to stage of infection (e.g., specimen obtained prior to the development of cellular immune responses).
- A Positive QFT should not be the sole or definitive basis for determining with M.tuberculosis. A Positive result should be followed by further medical evaluation and diagnostic evaluation for active tuberculosis disease (such as chest radiograph) are needed to exclude TB disease and confirm the diagnosis of LTBI.

This Test has

been Performed on QuantiFeron-TB Gold (QFT) ELISA test kit (FDA Approved)

Comments:

QuantiFeron TB Gold (Interferon Gamma Releasing Assay) test is whole blood test for detection of infection to *Mycobacterium tuberculosis* as occurs in active tuberculosis and latent tuberculosis infection (LTBI). If not detected and treated, LTBI may later develop into TB disease. This test measures the patient's immune reactivity to *M. tuberculosis*, the bacterium that causes TB. Blood samples are mixed with TB specific antigens and incubated for 20 to 24 hours. The antigens include ESAT-6 and CFP-10, proteins specific to tuberculosis complex. These antigens are not found in BCG strains or atypical Mycobacteria. If the patient is infected with *M. tuberculosis*, the patient's lymphocytes will recognize the antigens and release interferon -gamma in response.

Note: IGRA Test is approved as an in vitro diagnostic aid for detection of *Mycobacterium tuberculosis* infection (active disease and LTBI) and is intended for use in conjunction with risk assessment, radiography and other medical and diagnostic evaluations. The IGRA test does not differentiate between active and latent TB so latent patient will also be picked by IGRA. IGRA cannot be used as standalone test to diagnose TB infection. IGRA test is not established for any prognostic use.

Disclaimer : It cannot differentiate between latent infection and active tuberculosis.

*** End Of Report ***



Page 1 of 1


**DR. RUTURAJ MANIKLAL KOLHAPURE
MD, MICROBIOLOGIST**