





























<b>Name</b>	: Master. RUDRANSH	<b>Barcode No</b>	: B3425663
<b>Age /Gender</b>	: 2 YRS 8 MONTHS/Male	<b>Reg. No</b>	: 0012511110467
<b>Referring by</b>	:	<b>SPP Code</b>	: SPL-STS-032
<b>Referring Customer</b>	:	<b>Collected On</b>	: 11-Nov-25
<b>Primary Sample</b>	: Whole Blood	<b>Received On</b>	: 11-Nov-25
<b>Sample Tested In</b>	: Serum	<b>Reported On</b>	: 12-Nov-25

























<b>Test</b>	<b>Result</b>	<b>Units</b>	<b>Biological Ref.ranges</b>	<b>Method</b>	
Total IgE :	<b>275.4</b>	IU/mL	Age (Year) 0-<7 7-<19 (20-60) Adult	IU/mL <25-440 <25-450 0-160	CLIA

- Elevated Total IgE is observed in only 30% of patients with allergic rhinitis, 60% of patients with asthma and in 80 - 90% of patients with significant atopic eczema. It can also be elevated in 10 - 20% of patients with non-allergic rhinitis or non-allergic asthma, or other conditions such as allergic bronchopulmonary aspergillosis, some forms of immunodeficiency, neoplasia such as lymphoma, and parasitic disease. The measurement of Total IgE is the sum total of multiple individual allergen specific IgE levels.
- Atopic allergy implies a familial tendency to manifest conditions like Asthma, Rhinitis, Urticaria and Eczematous dermatitis either alone or in association with the presence of IgE.

## ALLERGY INHALANTS REPORT

Name of the Allergen	Result	Name of the Allergen	Result	Name of the Allergen	Result	Name of the Allergen	Result
Blomia tropicalis 	0.22	Dermatophagoide s microceras 	0.15	Dermatophagoides farinae 	0.20	Dermatophagoides pteronyssinus 	0.31
Cat dander 	0.29	Chicken feathers 	0.12	Cow dander 	0.21	Dog dander 	0.23
Duck feathers 	0.25	Finch feathers 	0.18	Goat Dander 	0.20	Horse Dandruff 	0.22
Guinea pig dandruff 	0.23	Parrot feathers 	0.21	Pigeon droppings 	0.13	Pigeon feathers 	0.25
Sweet vernal grass 	0.19	Barley grass 	0.20	Cocksfoot 	0.16	Common reed 	0.22
Cultivated oat 	0.22	Maize grass 	0.18	Ryegrass 	0.29	Cotton Dust 	0.21
House dust 	0.62	Jute Dust 	0.21	Rice dust 	0.16	Straw Dust 	0.18

Normal Range / Cut off for all allergens is : 0.35 U/L

Name of the Allergen	Result	Name of the Allergen	Result	Name of the Allergen	Result	Name of the Allergen	Result
Beet weed 	0.12	Common pigweed 	0.18	False ragweed 	0.29	Mugwort 	0.19
Ox eye daisy, Marguerite 	0.20	Red clover grass 	0.29	Western ragweed 	0.14	Timothy 	0.12
Jhonson grass 	0.23	Alternaria alternata 	0.21	Aspergillus fumigatus 	0.20	Aspergillus niger 	0.28
Candida albicans 	0.25	Cladosporium herbarum 	0.19	Mucor racemosus 	0.10	Penicillium notatum 	0.12
Trichoderma viride 	0.21	Cockroach 	0.25	Ant 	0.23	Honey Bee 	0.20
House Fly 	0.29	Wasp 	0.21	Moth 	0.20	House Cricket 	0.23
Body lotions	0.25	Deodorant	0.21	Hair Dye	0.21	Lip stick	0.20
Nail Polish	0.12	Perfume	0.24	Powder	0.25	Shampoo	0.28
Soap	0.10	Sun Creams	0.15	Latex	0.29	Paints	0.29
Papain	0.19	Plaster of Paris	0.28	Plastic	0.12	Silk	0.12
Smoke	0.23	Alcohol	0.29	Ash	0.18	Bone Cement	0.20
Detergent	0.29	Leather	0.21	Limestone	0.20	Nickel	0.26
Nylon fibers	0.21	Polyster	0.29	Sulfer	0.23	Wool mix	0.23
Yarn fibre	0.25	Cotton seed	0.23				

**Normal Range / Cut off for all allergens is : 0.35 U/L**

**Method: ELISA**

False positives occur in the following scenarios:

- *You have a small amount of IgE antibody to allergen but are not truly allergic to that. You can contact the allergen and experience absolutely no reaction to it.*

**Interpretation**

- Substances that cause an allergic reaction are called allergens. Besides dust and pollen, other common allergens include animal dander, foods, including nuts and shellfish, and certain medicines, such as penicillin.
- Allergy symptoms can range from sneezing and a stuffy nose to a life-threatening complication called anaphylactic shock. Allergy blood tests measure the amount of IgE antibodies in the blood. A small amount of IgE antibodies is normal. A larger amount of IgE may mean you have an allergy.